

## Cosmic Background Radiation temperature

Prof. A.K.T. Assis (*Apeiron*, 12, 10) studies the conservation of energy in a static Universe and the cosmic background radiation (CBR). He supposes that "the mean temperature of matter in the Universe is 2.7°K."

I have also looked into these topics, but I obtain a mean temperature for intergalactic matter ( $T_{av}$  of 5.4°K (see *Problems in the Study of the Universe*, 14, Leningrad, 1990, p. 210, in Russian), as follows:

$$T_F = \frac{T_{av}}{1+z_F} = \frac{T_{av}H}{c} \int_0^\infty \exp\left(-\frac{2H}{c}r\right) dr = \frac{T_{av}}{2}$$

where  $T_F$  and  $z_F$  are the temperature and cosmological redshift of the CBR, respectively,  $H$  is Hubble's constant and  $c$  is the velocity of light.

This matter may consist of hydrogen "snowballs" with a mass density of 0.076 gr cm<sup>-3</sup> and mean mass of 2000 gr. These would radiate energy in the form of transverse photons and absorb transverse (ordinary) photons and longitudinal photons (de Broglie 1943, 1949).

## References

- de Broglie, L. 1943, *Théorie générale des particules à spin*, Paris, p. 191.  
de Broglie, L. 1949, *Mécanique ondulatoire du photon et théorie quantique des champs*, Paris, Gauthier-Villars, chap. V.

Alexey Shlenov  
Budapeshskaya 66-1-77  
192286 St. Petersburg  
Russian Republic

## Cosmic Background Radiation: Reply to Shlenov

Dr. Shlenov discusses our work (Assis 1992) and points out that he has obtained a mean temperature for intergalactic matter of 5.4°K instead of 2.7°K. His model is an interesting one and deserves to be further investigated.

Here we would like to mention two other important references related to this subject in which the authors correctly estimated this temperature, prior to the experimental discovery by Penzias and Wilson in 1965. Finlay-Freundlich (1954) developed an alternative interpretation of the cosmological redshift on the basis of a tired light model, and as a corollary predicted the mean temperature

of intergalactic space between 1.9 and 6.0°K. Even earlier, Eddington, in a book first published in 1926, estimated the temperature of interstellar space at 3°K (Eddington 1988). And he obtained this with simple thermodynamic reasoning by equating the energy-density of the night sky with the Stefan-Boltzmann law according to which this density is proportional to the fourth power of the temperature.

## References

- Assis, A.K.T. 1992. "On Hubble's Law of Redshift, Olbers' Paradox and the Cosmic Background Radiation". in *Apeiron*, 12, 10-16.  
Eddington, A.S. 1988. *The Internal Constitution of the Stars*, Cambridge, Cambridge Univ. Press, Ch. 13, p. 371. Reprint of 1926 edition.  
Finlay-Freundlich, E. 1954. "Red Shifts in the Spectrum of Celestial Bodies". in *Philosophical Magazine*, 45, 303-319.

André K. T. Assis  
Department of Cosmic Rays and Chronology  
Institute of Physics  
State University of Campinas  
C. P. 6165, 13081 Campinas, S.P., Brazil

## Red-blue symmetry and action-at-a-distance

Ghosh (*Apeiron*, 12, 1) states in the first paragraph of his first letter: "thus, when photons graze massive objects...". His logic eludes me, in that if "blueshift during approach is neutralized by the equivalent amount of redshift when the photon recedes," he describes a canceled redshift. From whence does his observed redshift originate? How can the two shifts ever come together?

As I see it, observers, when they are in front of the approaching light, can see only a blue shift from behind a receding light. A redshift, in following a blue shift, cannot cancel the blue shift because they are separated in time, as well as by the light that makes the blue and red shifts.

Joop Nieland (*Apeiron*, 12, 2) states, "I am not bogged by action-at-a-distance...".

Newton erred with this thought, because gravity (mutual attraction) cannot be action-at-a-distance. Rather, it is an aura that surrounds all active celestial bodies, each to its correct distance in all directions, constantly. Action-at-a-distance would require intelligent celestial bodies, with fantastically versatile sensory abilities as well as magic

memories, to hold constant every movement of every planet, star, comet, etc.

How else but with a gravific aura and its own motion could the sun control all the movements of all its planets and comets?

Gravity is not a "field", in Einstein's sense. There are many gravities, each based on a specific celestial body. Some gravities overlies others. The system is so organized that none interferes in any other's business, because of weightlessness in orbit. The entire scheme is created in such a manner that there is complete order and precedence of orbital control.

*William Dunning  
Hollow Road  
Clinton Corners, N.Y. 12514*

### Corrections

APEIRON Number 8, Page 2: Equations (3) should read:

$$x' = (x - vt) \left( 1 - \frac{v^2}{c^2} \right)^{-1/2}$$

$$y' = y$$

$$z' = z$$

$$t' = \left( t - \frac{vx}{c^2} \right) \left( 1 - \frac{v^2}{c^2} \right)^{-1/2}$$

Equations (5) should read:

$$X' = X \cosh B - CT \sinh B \quad X = X' \cosh B + C'T' \sinh B$$

$$Y' = Y \quad Y = Y'$$

$$Z' = Z \quad Z = Z'$$

$$C'T' = CT \cosh B - X \sinh B \quad CT = C'T' \cosh B + X' \sinh B$$

Equations (9) should read:

$$X' = X \cosh B - CT \sinh B = X - VT$$

$$C'T' = CT \cosh B - X \sinh B$$

Page 3: The text before equation 16) should read: "...moving receivers whose trajectories pass through the source S, i.e. where..."

Page 5: In Figure 4, V7 should correspond to 8.00, not 6.00.

APEIRON Number 12 Page 3: Equation (9) should read:

$$n = 1 + \frac{2Zr_e}{r}$$

The paragraph before equation (11) should read: "Comparing equations (5) and (9) leads to the conclusion that  $\frac{c^2 r_e}{Gm_e} = \frac{e^2}{4\pi\epsilon_0 Gm_e^2}$  represents..."

The paragraph beginning with "Then why not...": The second sentence should end "...physically with rest mass somehow."

Page 12: Equation (3) should read:

$$F = \int_0^{\infty} \frac{Le^{-\alpha_L r}}{4\pi r^2} n 4\pi r^2 dr = \frac{Ln}{\alpha_L}$$

*The Editor*



## Note to authors

APEIRON is a scientific journal devoted to cosmology and fundamental physics. It publishes only convincingly argued, adequately documented papers in English, as well as English translations of historically important texts. APEIRON appears three times a year, and is indexed in *Physics Abstracts*. Submission implies that a paper is not under consideration by another publication and has not been published previously. All submissions will be reviewed by anonymous referees with expertise in the field, and every effort will be made to publish material presenting original viewpoints. There is no page charge or offprint charge. Send contributions to: C. Roy Keys, Editor, APEIRON, 4405 St-Dominique, Montreal, Quebec, Canada H2W 2B2.

All papers must be submitted typewritten or printed in three copies, and once accepted, on MS-DOS or MAC diskette (Word, RTF or ASCII format) or via Internet/ Compuserve whenever possible. The first page must include the title, the author's name and address, and a brief abstract. The International System is the preferred system of units. References must be indicated between brackets in the text by the author and year of publication, with a full alphabetical listing at the end of the manuscript. For books, the listing should include the full title, place of publication, publisher and year of publication. Journal articles should be referenced with the full title (between quotation marks), journal name, volume and page number. Graphical elements should be approximately 3x oversized, clearly numbered, labelled and captioned.